

Findings

32 J, 2016, D J, 27, 2015, J, 28, 2018, LA 32 ● 21,
 2016, D 24, 2017. E H 11454 2100
 LA ● 0 36 0 27 0 48), 20 7% (95% CI 14 6 26 7). A 2470
 , 2100 (85%) 2 2-
 LA, (● 0 39,
 0 24 0 65), 8 7% (3 5 14 0).
 H. (● 0 93, 0 74 1 16)
 (1 02, 0 73 1 43). LA IN \$316
 2 \$6518 2

Interpretation ● LA
 2 B D

We aimed to separately assess the effects of mHealth health messaging and PLA community mobilisation on the prevalence of intermediate hyperglycaemia and diabetes among the general adult population in rural Bangladesh, and to assess the effect of these interventions on the incidence of diabetes among people with inter-

and kitchen gardening to increase access to healthy food were popular strategies. Facilitators were locally recruited men and women who had completed higher secondary certificate level of education and were recruited by the study team following assessment of their communication skills, motivation, and familiarity with the study areas. Facilitators received a total of 14 days' training about PLA and community entry, group facilitation, and the basics of type 2 diabetes symptoms, prevention, and control. Each facilitator was responsible for running six to nine PLA groups each month. In addition, an equal number of men's and women's groups were established within each village, with a total of 122 groups facilitated by 16 facilitators (eight men, eight women) across 32 villages. Joint meetings of men's and women's groups were encouraged after phase 1 of the PLA cycle (ie, after identification and prioritisation of health determinants).

Training of informal health workers in the prevention and control of type 2 diabetes was done by the Diabetic Association of Bangladesh across all intervention and control villages during the intervention period. Project mapping of services in the study areas identified all informal care providers (eg, village doctors, pharmacy owners), who were then invited to participate in service-strengthening activities on a voluntary basis. This service strengthening included day-long workshops and provision of guidelines to cadres of largely unregulated care providers who had not received formal accredited training but might have had some degree of informal training through apprentices, workshops, or seminars. These informal care providers are typically the first point of care in health seeking by individuals in rural Bangladesh.²⁰

A sampling frame of all permanent residents aged 30 years and older was developed from a household census done between Aug 21, and Oct 28, 2015. 143 households with at least one eligible resident were then selected from each village by use of probability proportional to size sampling. A single eligible adult was selected from each of the 143 households for inclusion in the survey via simple random sampling. A baseline cross-sectional survey among the sampled individuals to obtain sociodemographic characteristics, behaviours, and knowledge of type 2 diabetes was done between Jan 23, and May 30, 2016. The survey included an overnight fasting blood glucose measurement in whole capillary blood obtained by finger prick in the middle or ring finger. All individuals without diagnosed type 2 diabetes then received a 75 g glucose load dissolved in 250 mL water. A 2-h post-prandial repeat capillary blood test was done to determine glucose tolerance status and differentiate between individuals with intermediate hyperglycaemia (defined as impaired fasting glucose or impaired glucose tolerance) and those with type 2 diabetes, based on WHO criteria (appendix).²¹ These baseline data were used to identify an intermediate hyperglycaemia cohort and to compare sociodemographic characteristics between the three trial groups.

Following intervention, the sampling frame was updated and a new random sample of adults aged 30 years and older was selected via the same sampling method as used at baseline. By chance, approximately 25% of individuals sampled for the end-of-study survey had also been included in the baseline survey. In addition, all individuals identified with intermediate hyperglycaemia at baseline were followed-up in the end-of-study survey to measure type 2 diabetes incidence in this cohort. An end-of-study survey of sociodemographic data, knowledge and behaviours, and anthropometric measures of weight, height, blood pressure, and fasting and 2 h post-prandial blood glucose measures was completed in the random cross-sectional sample and the baseline intermediate hyperglycaemia cohort between Jan 16, and April 30, 2018, using the same methods as at baseline.

Data were collected by 16 pairs of fieldworkers (one man and one woman in each pair) with at least secondary education, who underwent extensive training in survey methods, including supervised field practice. Most data collection took place in testing centres established by the field team for the purposes of the study, with additional data collection with pretested questionnaires taking place at respondents' homes.^{17,22} Data collectors were supervised by four field supervisors responsible for observing and verifying data. Data quality-control measures were implemented within the direct digital data capture system used (eg, range and consistency checks), through repeat measures by supervisors on a random basis, and where outlier data were detected on data inspection. Data collectors, supervisors, and managers were unaware of randomisation assignments at baseline, but might have been able to deduce assignment during data collection at the end of the study. Access to end-of-study data was restricted to the monitoring and evaluation managers until collection was complete, at which point the data were available for masked analysis by the lead author (EF).

Process evaluation data will be reported in detail elsewhere. We collected data in all four intervention upazillas of Faridpur district, including small group discussions with men and women attendees, with small groups of men and women with type 2 diabetes, and with group non-attendees. Later we met groups of men and women to explore triangulation and seek consensus on community changes. We also met with men's and women's group facilitators. In some of the groups we used participatory photography where they had identified and represented issues of importance to them using mobile phone cameras. Focus group discussions were digitally recorded and one author (KAK) made notes about the findings in English and translated field observation notes to English for analysis. Key themes around individual, household, and community change were compared with the theory of change drafted after the formative phase of research.

Outcomes

We prespecified two primary outcomes: the prevalence of intermediate hyperglycaemia and type 2 diabetes at the end of the study and the 2-year cumulative incidence of type 2 diabetes among the cohort with baseline intermediate hyperglycaemia.

Secondary outcomes were mean diastolic and systolic blood pressure, prevalence of hypertension, hypertension control (among those with known hypertension), BMI, prevalence of overweight and obesity and abdominal obesity (waist to hip ratio >0.9 for men and >0.85 for women), health related quality of life (using EQ-5D score), physical activity, fruit and vegetable consumption, and knowledge of the causes, symptoms, complications, prevention and control of type 2 diabetes. Additional secondary outcomes among people with type 2 diabetes were self-awareness of diabetes status and, among those with known diabetes, prevalence of diabetes control, psychological distress (with SRQ-20 screening tool), and receipt of professional medical treatment or advice for diabetes. Full specification of secondary outcomes and methods of assessment have been described previously.¹⁷

Additional pre-specified outcomes were intervention costs, incremental cost-effectiveness ratios and costs per disability adjusted life years (DALYs) averted for any effective intervention, and process indicators of intervention coverage and qualitative assessments of behaviour change caused by the interventions. An additional outcome of diabetes-only prevalence (excluding intermediate hyperglycaemia) was included post-hoc. Key findings from the qualitative, process assessments of behaviour change are summarised below, but full analyses are to be reported in a forthcoming publication.

Statistical analysis


We estimated that a target sample of 143 adults per village (total 13728, including 10% oversample for non-response) would provide 80% power at 95% confidence to detect a minimum 21.5% reduction in combined prevalence of type 2 diabetes and intermediate hyperglycaemia and 78% power to detect a 33% reduction in cumulative incidence of type 2 diabetes among the baseline intermediate hyperglycaemia cohort in intervention clusters relative to control clusters.¹⁷

Analysis of intervention effect on the combined prevalence of type 2 diabetes and intermediate hyperglycaemia included all individuals who provided blood

intermediate hyperglycaemia. Baseline characteristics were similar among all study groups (table 1, appendix), apart from a small difference in household wealth (higher in PLA villages), which was later adjusted for in

Increases in ability to report one or more valid causes, symptoms, complications, and strategies for prevention and control of diabetes were observed in both intervention groups compared with control, with the effect consistently greatest in the PLA group (table 4). Self-awareness of diabetes status among individuals identified as having type 2 diabetes by blood glucose testing was five-times higher in the PLA group than in the control group (aOR 5·09,

(table 3, figure 2). There was also no evidence of an effect of the mHealth intervention on the prevalence of type 2 diabetes alone in our post-hoc analysis (563 [15%] of 3797 vs 493 [13%] of 3821; adjusted [stratified, clustered design, and wealth] aOR 1·18, 0·95–1·48; $p=0\cdot139$).



was easier to initiate and sustain. PLA groups and their activities made community members feel in control of their health and able to prevent diabetes. The PLA intervention also destigmatised blood glucose testing and healthy behaviours such as physical activity, reduced or no sugar in tea, and healthy eating while socialising.

Total and average annual costs of the PLA intervention were INT\$601484 and \$240594, respectively. Total and average annual costs of the mHealth intervention were \$312630 and \$125052, respectively. The average annual costs of the PLA and mHealth per beneficiary (adults 30 years) covered were \$14 and \$7, respectively, with

costs per total population (all ages) being \$6 and \$3, respectively. The incremental cost-effectiveness ratios for PLA were \$316 per case of intermediate hyperglycaemia or type 2 diabetes prevented (or \$124 per DALY averted) and \$6518 per case of type 2 diabetes prevented (or \$2551 per DALY averted) among individuals with intermediate hyperglycaemia at baseline.

Discussion

We assessed two community interventions to prevent and control type 2 diabetes and intermediate hyperglycaemia in rural Bangladesh. Facilitated PLA community mobilisation led to large, significant reductions in the combined prevalence of type 2 diabetes and intermediate hyperglycaemia and 2-year incidence of type 2 diabetes among an intermediate hyperglycaemia cohort. The mHealth intervention had no effect on diabetes status. Both inter

activities and facilitated community engagement and intervention development activities. KAZ coordinated project activities in Bangladesh, co-led the project, and contributed to intervention development and study design. All authors contributed to the interpretation of study findings and had the opportunity to review and revise the final manuscript.

Declaration of interests

We declare no competing interests.

Data sharing

De-identified data collected for this study and a data dictionary are available from the corresponding author on reasonable request.

Acknowledgments

This work was funded by the UK Medical Research Council (MR/M016501/1) under the Global Alliance for Chronic Diseases Diabetes Programme. The study team thanks the DMagic trial steering committee (Graham Hitman [Chair], Martin McKee, Dina Balabanova, David Beran, Katherine Fielding, Lou Atkins, and Sophia Wilkinson), the data monitoring committee (Andrew Farmer [Chair], Margaret Thorogood, and Peter Byass) for their valuable contributions to the project. We are grateful to Fatima Zannat and Kazi Faisal Mahmud from mWorld for their contributions to the development of our mHealth intervention and Md Golam Azam for his contributions to field activities.

References

- 1 WHO. Global report on diabetes. Geneva: World Health Organization, 2016.
- 2 NIPORT, Mitra and Associates, ICF International. Bangladesh demographic and health survey 2011. Dhaka and Calverton: National Institute of Population Research and Training, Mitra and Associates, and ICF International, 2013.
- 3 Saquib N, Saquib J, Ahmed T, Khanam MA, Cullen MR. Cardiovascular diseases and type 2 diabetes in Bangladesh: a systematic review and meta-analysis of studies between 1995 and 2010. *BMC Public Health* 2012; 12: 434.
- 4 Bhowmik B, Afsana F, My Diep L, et al. Increasing prevalence of type 2 diabetes in a rural Bangladeshi population: a population based study for 10 years. *Diabetes Metab J* 2013; 37: 46–53.
- 5 Rahman MM, Akter S, Jung J, Rahman MS, Sultana P. Trend, projection, and appropriate body mass index cut-off point for diabetes and hypertension in Bangladesh. *Diabetes Res Clin Pract* 2017; 126: 43–53.
- 6 Fottrell E, Ahmed N, Shaha SK, et al. Diabetes knowledge and care practices among adults in rural Bangladesh: a cross-sectional survey. *BMJ Glob Health* 2018; 3: e000891.
- 7 Rawal LB, Tapp RJ, Williams ED, Chan C, Yasin S, Oldenburg B. Prevention of type 2 diabetes and its complications in developing countries: a review. *Int J Behav Med* 2012; 19: 121–33.
- 8 Ramachandran A, Snehalatha C, Ram J, et al. Effectiveness of mobile phone messaging in prevention of type 2 diabetes by lifestyle modification in men in India: a prospective, parallel-group, randomised controlled trial. *Lancet Diabetes Endocrinol* 2013; 1: 191–98.
- 9 Balagopal P, Kamalamma N, Patel TG, Misra R.