Appendix A. Notation 57

CHAPTER 1

Introduction

These notes are being prepared for a course in Fall 2017 in the Stanford Department of Mathematics. The ultimate goal of the course is to study the propagation of singularities on manifolds with boundary and parametrices for boundary value problems including the Melrose{Taylor parametrix. Throughout we will focus on the Dirichlet type problems, occasionally digressing to discuss the Dirichlet to Neuamnn map which can be applied to a wide variety of boundary conditions.

The material considered here comes largely from the work of Melrose [], Melrose{Sjostrand [], Taylor [], Zworski [] and Farris []

The basic example to which these notes apply is that of the wave operator $P = @_t^2 - _g$ posed on $R \quad M$ where M is a compact manifold with boundary $@M$

1. Notation for manifolds with boundary

2. Supported and Extendible distributions

We say that $a \in S^m(T^* M)$ if in any coordinates $(x; \ ) \ a(x; \ ) \in S^m(R^d \ R^d)$. Similarly, we say $a \in S^m_{phg}(T^* M)$ if $a(x; \ ) \in S^m_{phg}(R^d \ R^d)$. Then there is a a symbol map $_m : \ ^m_h(M) \ !$ $S^m(T^* M)=hS^{m \ 1}(T^* M)$ given by the following procedure. Let $(U \ ; \ )$ be an atlas on M. Then for $' \ ^2 \in C^1_c \ (U \ )$ a partition of unity on M,

$$(A) = \sum \ \sim \ ((\ ^1) \ ' \ A' \ )$$

where $\sim \ : T^* U \ ! \ T^* R^d$ is the lift of as a symplectomorphism. Note that $_m$ has the following properties

(1) $0 \ ! \ h \ ^m_h \ ^1 \ ! \ ^m_h \ ! \ ^m \ S^m=hS^{m \ 1} \ ! \ 0$ is exact.
(2) $(AB) = (A) \ (B)$.
(3) $([A; B]) = \ if \ (A); \ (B)g$ where $f \ ; \ g$ is the poisson bracket.

CHAPTER 3

Basic estimates for hyperbolic equations on manifolds with boundary

1. Energy Estimates and Well Posedness

1.1. Estimates without a boundary. We will work in the case of second order operators, but the methods developed here apply equally well to higher order equations.

1.1.1. *First order operators.* We consider the problem

$$P_t := (D_t - \mathrm{Op}(a_t))u = f; \quad 0 < t < T; \qquad u\,|_{t=0} = u_0 \tag{3}$$

where

(i) $a_t(x;\ ) = a(t;x;\ )$ belongs to a bounded set in $S^1(R^d \quad R^d)$ for $0 \quad t \quad T$
(ii) $t \mapsto a_t$ is continuous with values in $C^1\ (R^d \quad R^d)$
(iii) $\mathrm{Im}\, a(t;x;\ ) \quad M, 0 \quad t \quad T.$

We start with an energy estimate

Lemma 1.1. *Let* $s \in R$. *Then for* $\in R$ *large enough and all* $u \in C^1([0;T];H^s(R^d)) \cap C^0([0;T];H^{s+1}(R^d))$ *and* $p \in [1;1]$

$$\frac{1}{2}\Big(\int_0^T \|e^{-t}u(t;\ )\|_{H^s}^p\,dt\Big)^{\frac{1}{p}} \quad \|u(0;\ )\|_{H^s} + 2\int_0^T e^{-t}\|P_t u\|_{H^s}dt. \tag{4}$$

Proof. Then consider $E(t) = e^{-2t}\|u(t)\|_{L^2}^2$.

$$\begin{aligned}
\partial_t E(t) &= 2\,\mathrm{Re}\langle \partial_t[e^{-t}u]; e^{-t}u\rangle \\
&= 2\,\mathrm{Re}\, e^{-2t}\langle \partial_t u; u\rangle \quad 2E(t) \\
&= \quad 2e^{-2t}\,\mathrm{Im}\langle D_t u; u\rangle \quad 2E(t) \\
&= \quad 2e^{-2t}\,\mathrm{Im}\langle P_t u; u\rangle \quad 2e^{-2t}\,\mathrm{Im}\langle \mathrm{Op}(a_t)u; u\rangle \quad 2E(t) \\
&\quad 2\|e^{-t}P_t u\|E^{1=2}(t) + 2(C \quad)E(t)
\end{aligned}$$

where in the last line we apply the sharp Garding inequality (1.3) together with (i), (iii), and the fact that $u(t) \in H^1$. Choosing C, then

$$\partial_t E(t) \quad 2\|e^{-t}P_t u\|E^{1=2}(t).$$

Integrating in time gives

$$\sup_{0 \quad t} E(\ ) \quad E(0) + 2\sup_{0 \quad t} E^{1=2}(t) \int_0^{\cdot} \|e^{-s}P_s u\|ds$$

So,

$$\sup_{0\le t\le T} E^{1-2\rho}(\tau)\Big\|\int_0^t e^{-sA}P_s u\,ds\Big\|^2 \le E(0)+\Big\|\int_0^t e^{-sA}P_s u\,ds\Big\|^2$$

and in particular,

$$e^{-Ct}\|u(t)\| \le \|u(0)\| + 2\int_0^t \|e^{-Cs}P_s u\|\,ds.$$

Then, taking $\tau > 2C$ large enough and using that $\|e^{-t/2}\|_{L^p(R_t)} \le 2$ for $\tau > 0$. So,

$$e^{-\tau t}\|u(t)\| \le e^{(C-\tau)t}\|u(0)\| + 2\int_0^t e^{-\tau s}\|P_s u\|e^{(C-\tau)(t-s)}\,ds.$$

and taking $\tau > 2C$, and integrating both sides in t,

$$\Big(\int_0^T e^{-\tau t}\|u(t)\|^p\,dt\Big)^{1/p} \le \|e^{-t/2}\|_{L^p}\|u(0)\| + 2\int_0^T\Big(\int_0^t e^{-\tau s}\|P_s u\|e^{(C-\tau)(t-s)}\,ds\Big)^p dt\Big)^{1/p}.$$

Applying Minkowski's inequality then gives

$$\Big(\int_0^T e^{-\tau t}\|u(t)\|^p\,dt\Big)^{1/p} \le \|e^{-t/2}\|_{L^p}\|u(0)\| + \|e^{-t/2}\|_{L^p}\int_0^T e^{-\tau s}\|P_s u\|\,ds$$

Then, since $\|e^{-t/2}\|_{L^p} \le (2/\tau)^{1/p}$, the lemma follows for $s = 0$.

To finish the proof, apply the $s = 0$ case to $\tilde{A}_t = \mathrm{Op}(\langle\xi\rangle^s)A_t\,\mathrm{Op}(\langle\xi\rangle^{-s})$ with $u = \mathrm{Op}(\langle\xi\rangle^s)u$.

Theorem 1.1 (Well-posedness for first order equations). *Let (i)-(iii) hold and* $s \in \mathbb{R}$. *Then for all* $f \in L^1((0,T);H^s(\mathbb{R}^d))$ *and* $u_0 \in H^s(\mathbb{R}^d)$, *there is a unique solutions* $u \in C([0,T];H^s(\mathbb{R}^d))$ *of* (3) *and* (4) *holds.*

Proof. We start with uniqueness. Suppose (3) holds with $f = 0$ and $u_0 = 0$. Then, since $u \in C([0,1];H^s(\mathbb{R}^d))$, $\mathrm{Op}(a_t)u \in C([0,T];H^{s-1}(\mathbb{R}^d))$ and hence $\partial_t u \in C([0,T];H^{s-1}(\mathbb{R}^d))$. That is, $u \in C^1([0,T];H^{s-1})$ and in particular, (4) implies that $u = 0$.

To show existence, we apply the energy estimate to the adjoint problem. Suppose $v \in C_c^1((-1,T)\times\mathbb{R}^d)$. Then, observe that by (4) applied with $t \mapsto T - t$ gives for any $r \in \mathbb{R}$,

$$\sup_{t\in[0,T]}\|v(t)\|_{H^r(\mathbb{R}^d)} \le C\int_0^T \|(D_t - \mathrm{Op}(a_t))v\|_{H^r(\mathbb{R}^d)}\,dt.$$

Thus, for $f \in L^1([0,T];H^r(\mathbb{R}^d))$, $u_0 \in H^r(\mathbb{R}^d)$,

$$\begin{aligned}\int_0^T \langle f(t),v(t)\rangle_{\mathbb{R}^d}\,dt - \langle u_0, v(0)\rangle &\le (\|f(t)\|_{L^1([0,T];H^r(\mathbb{R}^d))} + \|u_0\|_{H^r})\sup_{[0,T]}\|v(t)\|_{H^r(\mathbb{R}^d)}\\ &\le C(\|f(t)\|_{L^1([0,T];H^s(\mathbb{R}^d))} + \|u_0\|_{H^r})\\ &\quad\int_0^T \|(D_t - \mathrm{Op}(a_t))v\|_{H^r(\mathbb{R}^d)}\,dt.\end{aligned} \tag{5}$$

By the Hahn-Banach theorem there exists $u \in L^1([0,T];H^r(R^d))$ so that

$$\int_0^T \langle u,(D_t - Op(a_t))v\rangle dt = \int_0^T \langle f(t),v(t)\rangle dt - i\langle h,v(0)\rangle$$

for all $v \in C_c^1((-\infty,T) \times R^d)$ in particular, u solves (3) as a distribution, so all we need to do is

with $P_j \in S^{2-j}$. Iterating this procedure gives

$$P = (D_t - \lambda_i)E_i + R$$

where $E_i = \sum_{k=0}^{l-1} E_{ik} D_t^k$ with $E_{ik} \in C^1(R; \Gamma^{l-1-k})$ and $R \in C^1(R; \Gamma^{2-l})$

Next, observe that

$$kD_t uk_{H^{s-1}} \quad k\ E_1 uk_{H^{s-1}} + k\ _2 uk_{H^{s-1}} + Ckuk_{H^{s-1}}$$

and hence

$$\sum_{k=0}^{1} kD_t^k uk_{H^{s+1-k}} \quad C \sum_i kE_i uk_{H^s} + Ckuk_{H^s}: \qquad {}_t uk_H \quad {}^k \tag{8}$$

Using (7), together with (4) applied to u

19 T0 T24

D^k

k=0

The Hahn Banach theorem implies that there exists $u \in L^1((0;T);H_{s+1})$ such that (10) holds with $Pu = f$, $D_t^j u(0) = u_j$.

We now need to show the other inclusions. Together with the estimates, they are obtained from the following lemma.

Lemma 2.2. *Let* $_{m;s}( ) = (h \ ^0i + i \ _1)^m h \ ^0i^s$: *Then* $_{m;s}(D)$ *is an isomorphism of* $S(\overline{R^d_+})$ *extending by continuity to* $L^2(R^d_+) \ ! \ H^{\ m; \ s}(\overline{R^d_+})$: *Moreover,* $_{m;s}(D)$ *extends by continuity from* $\overline{C^1_c}(R^d_+)$ *to an isomorphism from* $\overline{H}^{m;s}(R^d_+)$ *to* $L^2(R^d_+)$. *In particular,*

$$kuk_{m;s} = k \ _{m;s}(D)uk_{L^2(R^d_+)}$$

and $\overline{H}^{m;s}(R^d_+)$ *consists of those* $u \ 2 \ \overline{S}^0(R^d_+)$ *with* $_{m;s}u \ 2 \ L^2$.

Proof. One can easily check that $\operatorname{supp} F^{\ 1}(\ _{m;s}) \ \overline{R^d_+}$ and $\operatorname{supp} F^{\ 1}(\ _{m;s}) \ \overline{R^d}$: (For example by the Paley-Weiner theorem.) Therefore, $_{m;s}(D)$ maps $S(\overline{R^d}_+)$ continuously to itself and has inverse $_{m; \ s}(D)$. The extension to L^2 then follows since $j \ _{m;s}( )j^2 = h \ i^{2m} h \ ^0i^{2s}$: The second statement follows by duality.

We record the following consequence of Lemma 2.2 for later use.

Corollary 2.1. *Suppose* $u \ 2 \ H^{\ m; \ s}(\overline{R^d_+})$. *Then there exist* $u_0 \ 2 \ H^{1 \ m; \ s \ 1}(\overline{R^d_+})$ *and* $u_1 \ 2 \ H^{1 \ m; \ s}$ *so that*

$$u = u_0 + D_1 u_1:$$

Proof. Observe that $_{m;s}(D) = (hD \ ^0i + iD_1) \ _{m \ 1;s}$. So, since $_{m;s} : L^2 \ ! \ H^{\ m; \ s}$ is an isomorphism, there exists $v \ 2 \ L^2$ such that

$$u = \ _{m;s}v = (hD \ ^0i + iD_1) \ _{m \ 1;s}v$$

and observing that

$$hD \ ^0i \ _{m \ 1;s}v \ 2 \ H^{1 \ m; \ s \ 1}; \qquad i \ _{m \ 1;s}v \ 2 \ H^{1 \ m; \ s}:$$

2.0.2. *The local problem.* We start with the case $X = R^d$ and $(x) = x_1$. Then,

$$P = \sum_{j=0}^{2} P_j(x_1; x^0; D_{x^0}) D^j_{x_1}; \qquad P_j \ 2 \ Di^{\ 2 \ j}$$

and by the strict hyperbolicity assumption $0 < jP_2j$. Hence, dividing by a nonzero smooth function we may assume that $P_2 = 1$. Moreover, since P is a homogeneous polynomial of degree 2 in $(\ _1; \ ^0)$ for which that map $_1 \ 7! \ p(x; \ _1; \ ^0)$ has two distinct real zeros, we may assume P satis es the assumptions of Theorem 1.2 for $x \ 2 \ Y$.

Let $_1(x; \ ^0); \ _2(x; \ ^0)$ be the roots of $p(x; \ ; \ ^0)$. Fix $\ 2 \ C^1_c(R^d)$ with $\ 1$ in Y and de ne

$$p = \prod_{j=1}^{2} (\ _1 \ \tilde{} _j); \qquad \tilde{} _i = \ _j + (1 \ \ )j \ j \ ^0]:$$

Now, let $P' \in Di^{2}(R^d)$ with principal symbol p. Then P' satisfies the hypotheses of Theorem 1.2 for $x \in R^d$ and has constant coefficients outside a compact set.

Lemma 2.3. *Suppose that* P *is strictly hyperbolic with respect to* x_1 *in* $X \quad R^d$ *and let* $Y \quad X$ *be open and precompact. If* $f \in \overline{H}^{s;t}(R^d_+)$ *and* $s \quad 0$, $u_j \in H^{s+t+1-j}(R^{d-1})$, $j < 2$, *then there exists* $u \in \overline{H}^{s+1;t}(R^d_+)$ *such that* $D_1^j u \in C^0(R; H^{s+t+1-j}(R^{d-1}))$ *when* $x_1 \quad 0$ *and*

$$Pu = f \text{ in } Y \setminus R^d_+; \quad D_1^j u = u_j \text{ in } Y \setminus (R^{d-1} \quad f \quad 0g); j < 2:$$

Moreover, for $\in C_c^1$ (

Now, $U = u_0 + D_1 u_1 \in H^{s+1;t}$ and

$$P\tilde{U} - f = [P; D_1]u_1 \in H^{s+1;t-1}$$

since $[P, D$

Proof. Suppose that $Pu = 0$ in X_{0b}, $\operatorname{supp} u \quad f \quad (x) \quad 0g$. Then, by construction $\operatorname{supp} u$ $f > g$ where is uniform on X_{ab}. In particular, iterating nitely many times, $u = 0$.

For uniqueness with $s \quad 0$ an initial conditions imposed, suppose that $Pu = 0$ in X_{0b}, $u \; 2 \; \overline{H}^{1;t}(X_{0b})$ with $D^j u|_{t=0} = 0$ for $j < 2$. Then, extending u by 0 into < 0, we have $u \; 2 \; H^{1;t}(X$) we TJ/F40 10(j)]TJ/F41 7TJ/F41 7TJ/F41u 7TJ/F41 7TJ/F41 7TJ/o.t.2cF41 293(8[(H)]TJ/F33 7.8

4. Existence and Uniqueness

References: Hormande:III 24.1

Theorem 4.1. *Let* $f \in H^s_{loc}(X^o)$, $u_0 \in H^{s+1}(\partial X)$ *where* $s \geq 0$ *and assume* u_0, f *vanish on* $f < a$, *that* p *is strictly hyperbolic with respect to* $\quad$ *and* (11) *holds. Then there is a unique* $u \in H^{s+1}_{loc}(X^o)$ *such that*

$$\begin{cases} Pu = f & \text{in } X^o \\ u = u_0 & \text{on } \partial X \\ u = 0 & \text{on } \quad < a. \end{cases}$$

Moreover u *satisfies for every* $a' < a < b < b'$, [TODO]estimates

$$\|u\|_{H^{s+1}(a< <b\ )} \quad C(\|f\|_{H^s(a'< <b\ '')} + \|u_0\|_{H^{s+1}(\partial X)}).$$

CHAPTER 4

Propagation of Singularities

In this chapter we study an operator $P \in \mathrm{Diff}^2(X)$ for X a manifold with boundary ∂X and interior X^o. We assume throughout that ∂X is non-characteristic for P. That is, p does not vanish on $N^* \partial X$ For propagation in X^o, we will allow P to have an imaginary principal symbol. However, when it comes time to study the problem near the boundary, we will insist that the symbol be real valued.

1. Propagation in the bulk

2. Propagation of singularities for strictly hyperbolic problems

We now prove the propagation of singularities result for pseudodifferential operators.

Theorem 2.1. *Let X be a compact manifold and $P \in \Psi^m_{phg}(X)$ with $\sigma(P) = p - iq$ with p, q real valued. Suppose that $A, B, B_1 \in \Psi^0_{phg}(M)$ such that*

(1) for all $(x_0, \xi_0) \in \overline{WF}(A)$, there exist $T > 0$ so that $\exp(-Th$

is a di eomorphism onto its image and

$$(T \quad ; \quad T + ; V) \quad \text{ell}(B); \qquad (T \quad ; ; V) \quad `(B_1)$$

Let 0 (t) 2 C_c^1 (

We will study each of these regions separately. There is no propagation in E and the propagation in H results in broken bicharaceteristics. The propagation through G is subtle and will require a great deal of analysis.

4.1. The elliptic region. Our first task will be to prove the analog of the fact that if X is a compact manifold without boundary, then WF(u) WF(Pu) [f p = 0g. Note that the same proof shows that this continues to hold in the interior of a manifold with boundary and in

Also,

$$\|D_{x_1}u\|_{L^2}^2 \to \|D_{x_1}u\|_{L^2}^2,$$

completing the proof.

We now prove Lemma 4.2

Proof. We proceed using standard energy estimates.

$$(Pu,u)_{\overline{H}^{-1};H^1} = (D_{x_1}^2 u,u) + (b(x,D')D_{x_1}u,u) + (c(x,D')u,u)$$
$$\|D_{x_1}u\|_{L^2}^2 + (D_{x_1}u, b(x,D')u) + (c(x,D')u,u)$$

Now, since $T\{x_1 = 0\} \subset E$, there exists $\delta > 0$ so that

$$c(x,\xi') \geq \frac{b^2(x,\xi')}{4} + \delta|\xi'|^2.$$

Therefore, there exists $\delta_0 > 0$ so that

$$(c(x,D')u,u) \geq \frac{(1+\delta_0)}{4}\|b(x,D')u\|_{L^2}^2 - \frac{1}{2}\|r_{X'}u\|_{L^2}^2 - C\|u\|_{L^2}^2.$$

On the other hand

$$|(D_{x_1}u, b(x,D')u)| \leq \|D_{x_1}u\|_{L^2}\|b(x,D')u\|_{L^2} \leq \frac{1+\delta_0}{4}\|b(x,D')u\|_{L^2}^2 + \frac{1}{1+\delta_0}\|D_{x_1}u\|_{L^2}^2.$$

So,

$$|(Pu,u)| \geq c(\|D_{x_1}u\|_{L^2}^2 + \|r_{X'}u\|_{L^2}^2) - C\|u\|_{L^2}^2.$$

In particular,

$$c\|u\|_{H^1}^2 \leq \|Pu\|_{H^{-1}}\|u\|_{H^1} + (C+c)\|u\|_{L^2}^2 \leq \frac{1}{\epsilon}\|Pu\|_{H^{-1}}^2 + \epsilon\|u\|_{H^1}^2 + C\|u\|_{L^2}^2$$

So, choosing $0 < \epsilon < c/2$,

$$\|u\|_{H^1}^2 \leq C(\|Pu\|_{H^{-1}}^2 + \|u\|_{L^2}^2).$$

Next, let $E : H^s(\{x_1 = 0\}) \to H^s(\{x_1$

Lemma 4.4. *Suppose* $u \in L^2(R^d_+)$, $u|_{x_1=0} = u_0 \in H^{1/2}$ *and* $Pu \in H^{-1}$. *Then,*

$$\|u\|_{H^1} \leq C(\|Pu\|_{H^{-1}} + \|u_0\|_{H^{1/2}}).$$

Proof. Let $\chi \in C_c^1(R^{n-1})$ with $\chi \equiv 1$ near 0. Then put $u_\chi = \chi(D_\theta)u$. Then $u_\chi \to u \in L^2$, $Pu_\chi \to Pu \in H^{-1}$, $u_\chi \in H^{0,t}$ for all t. Now,

$$Pu_\chi = \chi(D_\theta)Pu + [\chi(D_\theta); b(x; D_\theta)]D_{x_1}u + [\chi(D_\theta); c(x; D_\theta)]u \in H^{-1,t}$$

for any t. In particular,

$$D_{x_1}^2 u_\chi \in H^{-1,t-1}; \qquad D_{x_1} u_\chi \in H^{-1,t}.$$

Therefore, $D_{x_1}u_\chi \in H^{0,t-1}$ and then, since $u_\chi \in H^{0,t-2}$, $u_\chi \in H^{1,t-2}$. In particular, choosing $t = 2$, $u_\chi \in H^1$. Hence,

$$\|u_\chi\|_{H^1} \leq C(\|Pu_\chi\|_{H^{-1}} + \|u_\chi\|_{L^2} + \|u_\chi|_{x_1=0}\|_{H^{1/2}}) \leq 2C(\|Pu\|_{H^{-1}} + \|u\|_{L^2} + \|u_0\|_{H^{1/2}}).$$

In particular, there exists a subsequence so that $u_\chi \rightharpoonup u$ in $\overline{H}^{-1}$. But $u_\chi \to u \in L^2$, so $u \in \overline{H}^{-1}$ and the estimate continues to hold.

Lemma 4.5. *Suppose* $s \geq 0$, $u \in \overline{H}^{s,t}(R^d_+)$, $u|_{x_1=0} = u_0 \in H^{s+t+\frac{1}{2}}$. *Then*

$$\|u\|_{\overline{H}^{s+1,t}} \leq C(\|Pu\|_{H^{s-1,t}} + \|u\|_{H^{s,t}} + \|u|_{x_1=0}\|_{H^{s+t+\frac{1}{2}}}).$$

Proof. We start with $t = 0$, Observe that

$$P(\langle D_\theta\rangle^s)u = \langle D_\theta\rangle^s Pu + [b(x; D_\theta); \langle D_\theta\rangle^s]D_{x_1}u + [c(x; D_\theta); \langle D_\theta\rangle^s]u$$

. So,

$$\|\langle D_\theta\rangle^s u\|_{H^1} \leq C(\|\langle D_\theta\rangle^s Pu\|_{H^{-1}} + \|u\|_{H^{0,s}} + \|\langle D_\theta\rangle^s u_0\|_{H^{1/2}}.$$

In particular,

$$\|u\|_{H^{1,s}} \leq C\|Pu\|_{H^{-1,s}} + \|u\|_{H^{s,0}} + \|u_0\|_{H^{s+\frac{1}{2}}}.$$

Iterating as before, we then obtain Therefore,

$$\|u\|_{H^{s+1,0}} \leq C\|Pu\|_{s-1,0} + \|u_0\|_{H^{\frac{1}{2}+s}}.$$

Now, for $t \neq 0$,

$$\|\langle D_\theta\rangle^t u\|_{H^{s+1,0}} \leq C(\|\langle D_\theta\rangle^t Pu\|_{H^{s-1,0}} + \|u\|_{H^{s,t}} + \|\langle D_\theta\rangle^t u_0\|_{H^{s+\frac{1}{2}}}$$

which concludes the proof.

Finally,

Lemma 4.6. *Suppose* $T \subset \{x_1 = 0\} \subset E$, $u \in N(\overline{R^d}_+)$,

$$Pu = f \in N(\overline{R^d}_+); \qquad u|_{x_1=0} = u_0.$$

Then,

$$WF_b(u)|_{T \setminus \{x_1=0\}} \subset WF_b(f)|_{\{x_1=0\}} \cup WF_b(u_0).$$

Proof. Here, the inclusion of the right hand side in the left is automatic from the de nition of WF_b. Therefore, we need only show that the right hand side is contained in the left. Suppose that $(0;\ _0^0) \not\in WF_b(f)|_{f x_1=0g} \cup WF_b(u_0)$. Fix $\ > 0$ so that

$$f0 < x_1 \quad g \setminus f\ x\ j\ (x;\ _1;0) \in WF(u) \cup WF(f)g = ;:$$

In particular, this is possible since $WF_b(u)|_{f x_1=0g} \quad T\ f x_1 = 0g$ and $WF_b(u)$ is closed. Therefore, for $x_1 > 0$ small and $(x;\ ) \in WF(u)$, $x_1 j\ _1 j < j\ ^0 j$.

Now, let

$$W^0 := WF_b(f)|_{f x_1=0g} \cup WF_b(u_0) \cup f\ (x;\ ^0)\ j\ 0 < x_1 \quad ;\ (x;\ ) \in WF(f) \text{ for some } _1 g:$$

This is closed since $WF_b(f)$ is. Let $\ \in S^1\ (R \quad T\ R^{d\ 1})$ with $\ (x;\ ^0) = 0$ for $x_1 \quad$ and order $1 \quad$ in a conic neighborhood of W^0. We call such a $\ $ a goodc cuto . Then, since $WF(f)$ avoids the $N\ f x_1 = cg$ for $c \quad ,\ (x; D^0)f\ \in C^1 \quad$ [TODO]references.

Now, choose $s;t$ such that for $\ \in S^j$ good,

$$(x; D^0)u \in \overline{H}^{s;t\ \ j}_{loc}:$$

Now, for $\ \in S^0$, good

$$P\ \ (x; D^0)u = \ \ (x; D^0)f\ + [P;\ \ (x; D^0)]u \in \overline{H}^{s\ \ 1;t}:$$

Indeed,

$$[P;\ \ (x; D^0)]u = P_0(x; D^0)\ \ _0(x; D^0) + D_{x_1} P_1(x; D^0)\ \ _1(x; D^0)$$

where $P_i\ \ _i$ is a good cuto in $S^{1\ \ i}$.

By iteration we obtain then that

$$(x; D^0)u \in \overline{H}^{s+1;t\ \ 1}$$

and in particular,

$$(x; D^0)u \in \overline{H}^{s_0;t_0}$$

with $s_0 \quad 0$.

Now, observe that

$$(x; D^0)u|_{x_1=0} = \ \ (x; D^0)u_0 \in C^1\ ;$$

So

$$P\ \ (x; D^0)u \in \overline{H}^{s_0\ \ 1;t_0};\qquad (x; D^0)u|_{x_1=0} \in C^1\ :$$

Therefore, by Lemma 4.5, $\ (x; D^0)u \in \overline{H}^{s_0+1;t_0}$ and in particular, $\ (x; D^0)u \in C^1$. Now, since $(0;\ _0^0) \not\in W^0$, we may take $\ (x;\ ^0)$ good with $\ (0;\ _0^0) = 1$ and hence by [TODO]reference, $(0;\ _0^0) \not\in WF_b(u)$.

Finally, we return to the general situation and prove our main theorem for singularities there.

Theorem 4.1. *Suppose that* $P \in Di\ ^2(X)$ *with* $@X$ *non-characteristic for* P. *Suppose* $f \in N(X)$ *and*

$$Pu = f \in X^o;\qquad u|_{@X} = u_0$$

Then,

$$WF_b(u)|_{@X} \setminus E\ = (WF_b(f)|_{@X} \cup WF(u_0)) \setminus E:$$

Definition 4.1. We de ne a *compressed broken bicharacteristic for* p as a continuous map

$$(t) : I \to \tilde{} \; n G$$

where $I \quad R$ is an interval and

- When $(t) 2 \tilde{} \setminus T X^{o}$, $(t) 2 C^{1}$ and $^{0}(t) = H_p((t))$:
- $f t 2 I j (t) 2 Hg$ is a discrete subset of I.

Locally near a point $(x^{0}; {}^{0}) 2 H$, such a broken bicharacteristic is given by

$$(t) = \begin{cases} (\exp(tH_p)(0; x^{0}; \sqrt{r_0(x^{0}; {}^{0})}; {}^{0})) & 0 \quad t \\ (\exp(tH_p)(0; x^{0}; -\sqrt{r_0(x^{0}; {}^{0})}; {}^{0})) & t \quad 0: \end{cases}$$

Our next theorem says that singularities of solutions to $Pu = 0$ with $uj_{@X} = 0$ are invariant along such broken bicharacteristics.

Theorem 4.3. *Suppose that* $P 2 Di^{2}(X)$ *with* $@X$ *non-characteristic for* P. *Suppose* $f 2 N(X)$ *and*

$$Pu = f 2 X^{o}; \qquad uj_{@X} = u_0$$

Suppose that $: I \to \tilde{}$ *is a broken bicharacteristic such that* $(I) \setminus (WF_b(f) [WF(u_0)) = ;$, *then for any* $t_0 2 I$ *if* $(t_0) \not\supseteq WF_b(u)$ *then*

$$(I) \not\supseteq WF_b(u):$$

Notice that for $(I) \setminus T @X = ;$, Theorem 4.3 is an easy consequence of Theorem 2.1. Therefore, in the proof we may work locally near a point $t 2 I$ such that $(t) 2 H$. Without loss of generality, we therefore assume $_0 = (0) 2 H$. Then, since $2 H$, freezing coe cients as in (14), we may replace P by P such that $T @X H$ so that Pu has the same wavefront set properties as Pu near ^{0} with $_0$

with $(_{j;}) = \sqrt{r(x; \ ^0)}$: We will show that the error can be improved to $E_{j+1} \ 2 \ S^{\ j}$. For this, let

$$_{j+1}(x; \ ^0) = \frac{(E_j)(x; \ ^0)}{2\sqrt{r(x; \ ^0)}} \ 2 \ S^{\ j}:$$

Then

$$\begin{aligned}
_{j+1}(x; D^0)(D_{x_1} \quad _{j;}) + E_j &= \ _{j+1}(x; D^0)(\ _{j;+} \quad _{j;}) + E_j + \ _{j+1}(x; D^0)(D_{x_1} \quad _{j;+}) \\
&= \ _{j+1}(x; D^0)(\ _{j;+} \quad _{j;}) + E_j \\
&\quad + (D_{x_1} \quad _{j;+}) \ _{j+1}(x; D^0) + [\ _{j+1}(x; D^0); D_{x_1} \quad _{j;+}] \\
&= E_{j+1;1}(x; D^0) \\
&\quad + (D_{x_1} \quad _{j;+} \quad _{j+1}(x; D^0)) \ _{j+1}(x; D^0) + E_{j+1;2}(x; D^0)
\end{aligned}$$

where $E_{j+1;1}; E_{j+1;2} \ 2 \ S^{\ j}$:

In particular,

$$\begin{aligned}
P &= (D_{x_1} \quad _{j;+})(D_{x_1} \quad _{j;}) + E_j(x; D^0) \\
&= (D_{x_1} \quad _{j;+})(D_{x_1} \quad _{j;}) \quad _{j+1}(x; D^0)(D_{x_1} \quad _{j;}) \\
&\quad + (D_{x_1} \quad _{j;+} \quad _{j+1}(x; D^0)) \ _{j+1}(x; D^0) + E_{j+1;2}(x; D^0) \quad E_{j+1;1}(x; D^0) \\
&= (D_{x_1} \quad _{j;+} \quad _{j+1}(x; D^0))(D_{x_1} \quad _{j;} + \ _{j+1}(x; D^0)) + E_j(x; D^0)
\end{aligned}$$

with $E_j \ 2 \ S^{\ j}$. Let $_{j+1;} = \ _{j;} \quad _{j+1}(x; D^0)$. Then de ning $_{0;} \quad \sum_{j \ 1} \ _j$ gives the desired factorization.

Repeating the arguments starting with the $\tilde{}$ factorization completes the proof

Next, we construct an operator $Q(x; D^0)$ with desirable microlocalization properties such that $[D_{x_1} \quad _+; Q] \ 2 \ S^{\ 1}$. This will allow us to complete the proof of Theorem 4.3 by e ectively microlocalizating $Pu = f$ to a broken bicharacteristic.

Lemma 4.8. *Fix* $_0 \ 2 \ H$, w3>>.505 Td [(0)]T817(pro)-27(of)-417(of)-416(Theorem)-416(4.3)-416(b)27(7006)]

Proof. Fix $q_0 \in S^0(T^*X)$ with $\operatorname{supp} q_0 \subset U \cap T^*_{\partial X}X$ and $q_0(\rho_0) = 1$. Then, since ∂_{x_1} is transverse to $\{x_1 = 0\}$, there exists $q_0 \in C^1([0,\epsilon) \times T^*X)$ solving

$$(\partial_{x_1} - H^p -$$

Observe that

$$(D_{x_1} \tilde{\ }) Q (D_{x_1} \tilde{\ }_+) = Q P \tag{22}$$

implies $_0 \not\geq WF_b(u)$. Switching the roles of Q , we also obtain that if ( ) $\not\geq WF_b(u)$, then $_0 \not\geq WF_b(u)$ which completes the proof.

[TODO]state estimates

5. The generalized bicharacteristic ow

6. The Weyl law on a manifold with boundary

7. Microlocal defect measures

CHAPTER 5

Equivalence of Glancing Hypersurfaces

1. Symplectic preliminaries

2. Folding relations

3. The billiard ball maps

4. Formal solution

5. Completion of the proof

6. Some consequences

Therefore, it suffices to find an P even function u_0 on $Q \setminus P$ so that

$$[\exp(-r)u_0]_O = c[\exp(-r)u_0]_E + f, \qquad u_0(\ ) = c_0.$$

Moreover, since

$$[\ \ ]_E = [\ ]_E[\ ]_E + {}^2[\ ]_O[\ ]_O$$
$$[\ \ ]_O = [\ ]_O[\ ]_E + [\ ]_E[\ ]_O$$

and $[\exp(-r)]_E(\ ) \neq 0$, it is enough to find an P even function, v with

$$[v]_O = c[v]_E + f, \qquad v(\ ) = c_0.$$

for some given Q even c and f.

Lemma 4.1. *For any Q even $c, f \in C^1(Q \setminus P)$, $c_0 \in C$, there exists a smooth P even function v on $Q \setminus P$ so that*

$$[v]_O = c[v]_E + f, \qquad v(\ ) = c_0 \tag{43}$$

in a neighborhood of .

Proof. For this, we work in the normal form so that

$$_Q(y_2, y'', \ _1, \ ^0) = (y_2, y'', \ \ _1, \ ^0), \qquad _P(y_2, y'', \ _1, \ ^0) = (y_2 \ \ 2\ _1, y'', \ \ _1, \ ^0).$$

In this coordinates, then we seek an P even function solving (43). To simplify notation we write $t_1 = \ _1$ and $t_2 = y_2$, $t' = (y'', \ ^0)$ so that

$$_Q(t_1, t_2, t'') = (\ t_1, t_2, t'), \qquad _P(t_1, t_2, t') = (\ t_1, t_2 \ \ 2t_1, t').$$

Since v is P even, $v = v(t_1^2, t_2 \ \ t_1, t')$. We start by solving for v

Hence, by linearity, we need only solve

$$[v]_O = c[v]_E + e; \qquad e \in C^1 ; \ {}_Q e = e; \quad @e(t) = O\ (t_1^1\ ); \quad v|_{t_1=0} = 0 \tag{46}$$

Since we are interested in solving near 0, we may also assume that supp e f $t_2 < 1$g, c is supported near $t_2 = 0$, and supp v f $t_2 < 1$g.

Now, observe that

$$\frac{v \quad {}_Q v}{2t_1} = c(t)\frac{v + {}_Q v}{2} + e(t)$$

implies

$$v = \frac{1}{1 \quad t_1 c(t)}[(1 + t_1 c(t))\ {}_Q v + 2t_1 c(t) e(t)]: \tag{47}$$

Since ${}_Q e = e$, ${}_Q c = c$ ${}_P v = v$, ${}_P t_1 = \ t_1$,

$$\begin{aligned} {}_P v &= \frac{1}{1 + \ {}_P t_1\ {}_P c(t)}[(1 + \ {}_P t_1\ {}_P c(t))\ {}_P\ {}_Q v + 2\ {}_P t_1\ {}_P c(t)\ {}_P \quad e(t)] \\ v &= \frac{1 \quad t_1\ {}_P c(t)}{1 \quad t_1\ {}_P c(t)}\ {}_P\ {}_Q v \quad \frac{2t_1\ {}_P c(t)}{1 \quad t_1\ {}_P c(t)}\ {}_P\ {}_Q e(t) \\ &= \frac{1 \quad t_1 \quad c(t)}{1 \quad t_1 \quad c(t)}\ \ v + \frac{2t_1 \quad c(t)}{1 \quad t_1 \quad c(t)} \quad e(t) \end{aligned} \tag{48}$$

where $\ = {}_P \quad {}_Q$ is the induced billiard ball map. Letting

$$a(t) := \frac{1 \quad t_1 \quad c(t)}{1 + t_1 \quad c(t)}; \qquad b(t) := \frac{2t_1 \quad c(t)}{1 \quad t_1 \quad c(t)};$$

we then have

$$v = \sum_{m=0}^{M} B_m(\ {}^{m+1})\ e + A_M(\ {}^{M+1})\ v \tag{49}$$

where

$$B_m = (\ {}^m)\ b \prod_{k=0}^{m-1} (\ {}^k)\ a; \qquad A_M = \prod_{k=0}^{M} (\ {}^m)\ a:$$

Now, note that

$$(t_1; t_2; t') = (t_1; t_2 + 2t_1; t')$$

and hence, since supp v; supp e f $t_2 < 1$g, for $t_1 > 0$, $t_2 > \ 1$, and $M > 2t_1^{\ 1}$, $(\ {}^M)\ v = (\ {}^M)\ e = 0$. In particular, the the right hand side of (49) is independent of $M > 2t_1^{\ 1}$ when t_1

We clearly have

$$|1 - b(t)| \le c|t_1|$$

and hence, for $m \le M = 2t_1^{-1}$,

$$|B_m(t)| \le (1 + c|t_1|)^M < C < 1 :$$

Therefore, the coe cients on (∂^{m+1}) e in (50) are uniformly bounded. Now, e vanishes to all orders at $t_1 = 0$ and is supported in $t_2 < 1$. Therefore, for all $N > 0$, there exists $c_N > 0$ so that

$$|(\partial^m) g(t)| \le C_N |t_1|^{2N} (1 + |t_2| + 1 + m|t_1|)^{-N} \le c_N |t_1|^N (1 + m)^{-N} :$$

In particular, the sum (50) converges rapidly and hence de nes a continuous function vanishing to all orders at $t_1 = 0$.

Now, B_m is a product of at most M terms. Thus, when it is di erentiated, in $(t_2; t')$, it produces at most M choose N terms each of which is bounded by $c_N |t_1|$. In particular, di erentiation N times produces at most M^N sums of the form (50) except that sum numb16 10.9091 Tc (<)]TJ/F16 10.9091 T
[(1 997]TJ/F42 10.9091 Tf 8.069 1.689 Td [(j)]TJ/F40 10.9091 Tf 3.03 0 Td u

CHAPTER 7

Calderon Projectors { Boundary Integral Operators??

APPENDIX A

Notation

- $\mathrm{Di}^{m}(M)$ differential operators of order m on a manifold M.