AVERAGES OF EIGENFUNCTIONS OVER HYPERSURFACES

YAIZA CANZANI, JEFFREY GALKOWSKI, AND JOHN A. TOTH

Abstract. Let $(M;g)$ be a compact, smooth, Riemannian manifold and $f_h g$ an L^2-normalized sequence of Laplace eigenfunctions with defect measure . Let H be a smooth hypersurface with unit exterior normal : Our main result says that when is *not* concentrated conormally to H, the eigenfunction restrictions to H satisfy

$$\int_H {}_h d_H = o(1) \quad \text{and} \quad \int_H hD \; {}_h d_H = o(1);$$

$h \to 0^+$.

1. Introduction

On a compact Riemannian manifold $(M;g)$, with no boundary, consider a sequence of Laplace eigenfunctions $f_h g$,

$$h^2 \; g \; {}_h = {}_h;$$

normalized so that $k_h k_{L^2(M)} = 1$. The goal of this article is to study the average oscillatory behavior of ${}_h$ when restricted to a hypersurface $H \subset M$. Namely, the goal is to find a condition on the pair $(f_h g; H)$ that so that

$$\int_H {}_h d_h = o(1); \tag{1}$$

as $h \to 0^+$, where ${}_H$ denotes the hypersurface measure on H induced by the Riemannian structure.

It is important to point out that one cannot always expect to observe this oscillatory decay. For instance, on the round sphere, zonal harmonics of even degree integrate to a constant along the equator. Also, for any closed geodesic inside the square at torus. there is a sequence of eigenfunctions that integrate to a non-zero constant.

Integrals of the form (1) have been studied for quite some time, going back to the work of Good [Goo83] and Hejhal [Hej82] that treated the case where H is a periodic geodesic inside a compact hyperbolic manifold. These authors proved that in such a case, $\int_H {}_h d_h = O(1)$ as $h \to 0^+$. This result was generalized by Zelditch [Zel92] to the case where H is a hypersurface inside a compact manifold. Indeed, it follows from [Zel92] that for a density one subsequence of eigenvalues $fh_j g_j$; one has $\lim_{j\to\infty} \int_H {}_{h_j} d_H = 0$: Moreover, one can actually get an explicit polynomial bound of the form $O(h^{1/2 \; 0})$ for the rate of decay of expectations for the density-one subsequence (see [JZ16]). However, the latter estimate is not satisfied for all eigenfunctions and it is not clear which sequence of eigenfunctions must be removed for the estimate to hold. There are several articles that address this issue by restricting to special cases of

Riemannian surfaces $(M;g)$ and special curves $H \subset M$: When $(M;g)$ is a compact surface with constant curvature equal to -1 and H is a closed geodesic Reznikov [Rez15] proved that the periods $\int_H h\, d\ell_h = O(1)$. Working on surfaces of strictly negative curvature, and choosing H to be a geodesic, Chen-Sogge [CS15] improved this result to a o

2. Decomposition of defect measures

2.1. **Invariant Measures near transverse submanifolds.** Let N be a smooth manifold, V be a vector field on N and write $\varphi_t^V : N \to N$ for the flow map generated by V at time t. Let $\mathcal{N}$ be a smooth manifold transverse to V. Then for $\varepsilon > 0$ small enough, the map $\Phi : (-2\varepsilon, 2\varepsilon) \times \mathcal{N} \to N$

$$\Phi(t, q) = \varphi_t^V(q)$$

is a diffeomorphism onto its image and we may use $(-2\varepsilon, 2\varepsilon) \times \mathcal{N}$ as coordinates on N near $\mathcal{N}$.

Lemma 5. *Suppose that μ is a finite Borel measure on N and that $V \mu = 0$ i.e. $(\varphi_t^V)^* \mu = \mu$. Then, for a Borel set $A \subset [-\varepsilon, \varepsilon] \times \mathcal{N}$,*

$$\mu(A) = dt \, d\nu(A)$$

where ν is a finite Borel measure on $\mathcal{N}$.

Proof. Throughout this proof, we slightly abuse notation by identifying $\mathcal{N}$ with $\{0\} \times \mathcal{N}$. For $B \subset \mathcal{N}$ Borel, define

$$d\nu(B) :=:$$

Then, since

$$([a;b] \quad B) \quad ([\ '' + \tfrac{2''(k_1 \quad 1)}{n};\ \ '' + \tfrac{2''k_2}{n}] \quad B) = \sum_{j=k_1}^{k_2} ([\ '' + \tfrac{2''(j \quad 1)}{n};\ \ '' + \tfrac{2''j}{n}] \quad B);$$

$$(I \quad B) \quad \frac{2''(k_2 \quad k_1 + 1)}{n} \frac{1}{2''} \ ([\ '';'') \quad B) \quad (b \quad a + 2\ )d \quad (B):$$

Sending $\ !\ 0$ proves

$$(I \quad B) \quad jIjd \quad (B):$$

Therefore, if $a = b$, $(I \quad B) = 0$ and we may assume $a < b$. Fix $\ > 0$ so that $b \quad a > 2\ > 0$ and choose

$$a \quad '' \quad 1 + \tfrac{2(k_1 \quad 1)}{n} \quad a + \quad b \quad '' \quad 1 + \tfrac{2k_2}{n} \quad b:$$

Then,

$$(I \quad B) \quad \frac{2''(k_2 \quad k_1 + 1)}{n} \frac{1}{2''} \ ([\ '';''') \quad B) \quad (b \quad a \quad 2\ )d \quad (B)$$

and sending $\ !\ 0$ proves (3) and hence nishes the proof of the lemma.

2.2. **Fermi coordinates.** Let $H \quad M$ be a closed smooth hypersurface and let U_H be a Fermi collar neighborhood of H. In Fermi coordinates

$$U_H = f(x^0; x_n) : x^0 2 H \text{ and } x_n 2 (\ c; c)g$$

for some $c > 0$, and $H = f(x^0; 0) : x^0 2 Hg$. Since H is a closed hypersurface, it divides M into two connected components $\ _H$ and $M n\ _H$. In the Fermi coordinates system, the point $(x^0; x_n)$ is identi ed with the point $\exp_{x^0}(x_n\ _n) \ 2 \ U_H$ where $\ _n$ is the unit normal vector to $\ _H$ with base point at $x^0 2 H$.

The Fermi coordinates on U_H induce coordinates $(x^0; x_n; \ ^0; \ _n)$ on $S_U M = f(x; \ ) 2$ 2 ($-4 Tf 65.104 0 T.

2.3. Transversals for defect measures. We now apply Lemma 5 to the special case of defect measures, using the fact that they are invariant under the geodesic flow. In what follows we write $|\xi|_{x^0} := |\xi|_{g_H(x^0)}$, where g_H is the Riemannian metric on H induced by g. Let

$$G_H(\lambda) := \{(x,\xi) \in S_H M : |\xi|^2_{x^0} = 1 = \lambda^2 g\},$$

and define the set of non-glancing directions

$$\Gamma := S_H M \cap G_H(\lambda).$$

Lemma 6. Suppose μ is a defect measure associated to a sequence of Laplace eigenfunctions. Then, for all $\Gamma > 0$ there exists $\varepsilon > 0$ small enough so that

$$\Phi^* \mu = dt\, d\nu \quad \text{on } (\Gamma - \varepsilon, \Gamma + \varepsilon)$$

where

$$\Phi : (\Gamma - \varepsilon, \Gamma + \varepsilon) \times \bigcup_{|s|<\varepsilon} G^s(\Gamma), \qquad (t,q) = G^t(q),$$

is a diffeomorphism and $d\nu$ is a finite Borel measure on Γ.

Proof. In what follows we use Lemma 5 with $N = S^*M$, $V = H_p$ the Hamiltonian flow for $p = |\xi|_g$, and $\varphi_t^V = G^t$ the geodesic flow. Note that since μ is a defect measure for a sequence of Laplace eigenfunctions, it is invariant under the geodesic flow G^t. Then, for q_1 Tf 4.814 1.636 Td 10.612 0 f 11.772 0 0 Td [(091 Tf 10.225 0 Td [( )]TJ/F -3.959 Td [(M)]TJ/F15 10.